



Review of grouper hatchery technology

by Mike Rimmer¹

The following article is a slightly condensed version of an article published in the *Grouper Aquaculture Electronic Newsletter*, 27 January 2000. (<http://naca.fisheries.go.th>)

Larval rearing of groupers

Successful larviculture of groupers has been constrained by generally poor and irregular survival. The principal constraints to successful larviculture (Kohno *et al.*, 1990 and 1997; Tamaru *et al.*, 1995; Leong, 1998; Rimmer *et al.*, in press) are:

1. the small gape of the larvae and hence their requirement for small prey at first feed; and
2. the occurrence of high mortality at various stages through the larval rearing phase.

This document briefly reviews grouper larviculture technology, and summarises the current status of this technology.

Taxonomic note

Because of the sometimes-confused taxonomic status of groupers, particularly the genus *Epinephelus*, the literature pertaining to grouper aquaculture commonly misidentifies the species concerned. There is a voluminous literature on *E. tauvina*, but most of this in fact refers to the estuary cod (or greasy grouper) *E. coioides*. To add to the confusion, the synonym *E. suillus* is also sometimes used to refer to *E. coioides*. The black-spotted cod *E. malabaricus* is sometimes referred to by its synonym *E. salmoides*. However, much of the Thai literature on *E. malabaricus* apparently refers to *E. coioides* (R. Yashiro, pers. comm.). In the following literature review, references to *E. tauvina* and *E. suillus* are assumed to refer to *E. coioides*, and references to *E. salmoides* to *E. malabaricus*. Instances of possible misidentification of *E. coioides* as *E. malabaricus* are noted.

Status of grouper hatchery technology

Eggs of *E. coioides*, *E. fuscoguttatus*, *Cromileptes altivelis* and *Plectropomus leopardus* take 15–19 hours to hatch, while the European white grouper, *E. aeneus* take around 25 hours to hatch (Rimmer *et al.*, in press).

Grouper eggs and newly hatched larvae are very sensitive to stress and handling (Predalumpaburt & Tanvilai, 1988; Caberoy & Quinitio, 1998). Handling mortality is minimised by handling only neurula-stage eggs (after the formation of the optic vesicles) and by stocking eggs into the culture tanks 2 hours before hatching so that the larvae need not be handled (Lim, 1993; Tamaru *et al.*, 1995; Caberoy & Quinitio, 1998). Grouper larvae are stocked at relatively high density: 20–30 per litre (Ruangpanit, 1993; Duray *et al.*, 1996) up to 50 per litre (Lim *et al.*, 1986; Aslianti, 1996). The larvae are sensitive to light during the early stages of their development and are generally kept in darkened conditions.

Rearing tanks are generally rectangular and range in size from 5 to 30 m³ (Rimmer *et al.*, in press). Tank size, shape and colour may affect the survival of grouper larvae cultured intensively. *E. coioides* larvae cultured in 3 m³ tanks demonstrated a better survival rate (19.8%) at Day (D) 24 compared with only 7.4% for those in 0.5 m³ tanks at D21 (Duray *et al.*, 1997). Growth and survival of *E. fuscoguttatus* larvae was improved using cylindrical rather than rectangular larval rearing tanks (Waspada *et al.*, 1991b). *Cromileptes altivelis* larvae exhibited higher survival in green or blue coloured tanks than in red or yellow coloured tanks, but growth rate was not affected by tank colour (Aslianti *et al.*, 1998). Grouper larval rearing tanks are usually supplied with microalgae (generally *Nannochloropsis oculata*, formerly known as marine *Chlorella*), or *Tetraselmis* sp. at 500 x 10³ to 100–500 x 10⁶ cells per ml ('green water' system) (Ruangpanit, 1993; Tamaru *et al.*, 1995; Watanabe *et al.*, 1996; Leong, 1998; Rimmer, 1998; Rimmer *et al.*, 1998). The microalgae provides a shading effect, provides food for the live prey organisms added to the tanks, and may also be ingested by the larvae (although whether the larvae gain any nutritive value from ingested microalgal cells is unknown). More recently, the use of *Isochrysis* for 'green-water' rearing has been shown to improve larval growth and survival (Su *et al.*, 1998).

1. Queensland Department of Primary Industries, Northern Fisheries Centre, Cairns, Queensland, Australia

The mouth of larval groupers generally opens 2–3 days after hatching (D2–3), and the larvae begin feeding soon thereafter (Kitajima *et al.*, 1991; Kungvankij *et al.*, 1986; Ruangpanit, 1993; Ruangpanit *et al.*, 1993; Duray 1994; Doi *et al.*, 1997). Kohno *et al.* (1997) described the development of the feeding apparatus in *E. coioides* in detail and suggested that the major difficulties in larval rearing of groupers were attributable to the small size of the bony elements forming the oral cavity, small mouth and body size, poor reserves of endogenous nutrition and lower initial feeding rates.

Grouper larvae are initially fed on rotifers, often in combination with oyster trochophores, mussel larvae, sea urchin eggs or barnacle nauplii (Hussain & Higuchi, 1980; Lim, 1993; Ruangpanit, 1993; Tamaru *et al.*, 1995; Watanabe *et al.*, 1996; Rimmer *et al.*, 1998). Oyster trochophores, mussel larvae, sea urchin eggs and barnacle nauplii are around 70 µm in size and are thus small enough to be readily consumed by the larvae (Kungvankij *et al.*, 1986; Tamaru *et al.*, 1995). Small-strain (S-type) rotifers (*Brachionus rotundiformis*) are too large for newly hatched grouper larvae to ingest. Super-small-strain (SS-type) rotifers (*Brachionus* sp.), or S-type rotifers screened to <90 µm, are suitable for grouper larvae at first feed (Lim, 1993; Tamaru *et al.*, 1995; Watanabe *et al.*, 1996; Duray *et al.*, 1997; Su *et al.*, 1997). Optimal prey density during the early larval stages is 10–20 organisms/ml (Ruangpanit, 1993; Ruangpanit *et al.*, 1993; Tamaru *et al.*, 1995; Wardoyo *et al.*, 1997).

The use of copepod nauplii as prey during the early larval rearing of groupers has shown considerable promise in improving larval growth and survival (Hussain & Higuchi, 1980; Doi *et al.*, 1997). Provision of nauplii of calanoid copepods (mainly *Pseudodiaptomus annandalei* and *Acartia tsuensis*) and rotifers in rearing tanks, resulted in higher survival and faster growth in *E. coioides* compared with the use of rotifers alone (Doi *et al.*, 1997; Toledo *et al.*, 1999). *E. coioides* larvae actively select copepod nauplii in preference to rotifers (Toledo *et al.*, 1997).

S-type rotifers are fed from about D7 (Tamaru *et al.*, 1995; Su *et al.*, 1997), and brine shrimp (*Artemia franciscana*) are introduced from about D10. Brine shrimp are initially fed at 1–3/ml, increasing gradually to 7–10/ml at D25–35 (Ruangpanit, 1993). Higher prey densities (2–3/ml) enhance growth and survival (Duray *et al.* 1997), as does frequent feeding of brine shrimp (4–5 times per day) (Ruangpanit, 1993; Duray *et al.*, 1997). Grouper larvae fed rotifers and brine shrimp enriched with *n*-3 HUFAs demonstrate better growth, survival and greater stress endurance than those fed with normal live feeds (Pechmanee *et al.*, 1988; 1993;

Pechmanee & Assavaaree, 1993; Chao & Lim, 1991; Dhert *et al.*, 1991; Lim, 1993; Qunitio, 1996). Most laboratories supplement live prey for grouper larvae with microalgae (e.g. *Tetraselmis*, *Chaetoceros*), home-made lipid emulsions (egg yolk blended with cod liver oil), or commercially available lipid emulsions or microencapsulated supplements (Ruangpanit 1993; Rimmer *et al.*, in press). Minced fish or shrimp may be introduced from about D35 to wean the larvae (or juveniles) from live to inert feeds (Hussain & Higuchi, 1980; Ruangpanit, 1993; Tamaru *et al.*, 1995).

Survival of groupers to metamorphosis is generally low: usually <10% and commonly <1%. Recent reports of survival in experimental conditions are: 1–10% (average 4%) for *E. coioides* (J. Toledo, pers. comm.) and 1–5% for *C. altivelis* (K. Sugama, pers. comm.). In addition to low average larval survival, survival is highly variable, and a (relatively) successful larval rearing 'run' can be followed by several runs which have negligible or zero survival. It is this both these aspects, low and irregular survival, that have constrained the application of the existing fingerling production technology to commercial hatchery production.

The major causes of mortality in grouper larviculture are:

1. High mortality associated with the commencement of exogenous feeding. This mortality may be associated with the provision of live prey organisms of unsuitable size and nutritional composition, but even when 'suitable' prey types are used, there is generally high mortality at this stage (Ordonio-Aguilar *et al.*, 1995; Duray *et al.*, 1997).
2. Several mortality syndromes have been described for grouper larvae. A commonly reported mortality syndrome is the 'shock syndrome' that occurs in late stage larvae from about D25 (Lim, 1993; Duray *et al.*, 1997). This problem may be related to nutritional deficiencies in the live prey organisms used to feed the larvae, since shock syndrome is symptomatic of low levels of HUFAs in the diet (Cowey & Sargent, 1972).
3. Cannibalism is a major cause of mortality during the later stages of larval rearing, i.e. from D30–35 (Lim, 1993; Tamaru *et al.*, 1995; Rimmer *et al.*, in press). Although cannibalism can be controlled by grading larvae and juveniles into similar size classes, grading is often associated with high mortality because handling commonly induces the shock syndrome seen in late stage larvae (Rimmer *et al.*, in press). Provision of shelter is reported to reduce cannibalism in juvenile *E. malabaricus* (*E. coioides*?) (Rimmer *et al.*, in press).

Nutritional requirements of larval groupers

The few studies that have been carried out on the nutritional requirements of larval groupers suggest that supplementation of the larval diet with HUFAs improves growth and survival. Waspada *et al.* (1991c) found higher growth rates of *E. fuscoguttatus* larvae when they were fed rotifers supplemented with bakers yeast with sardine oil or bakers yeast with cod oil. These treatments had high levels of EPA (8.8–8.9% respectively) but highly variable levels of DHA (0.1–5.5% respectively) (Waspada *et al.*, 1991a). Su *et al.* (1997) found that growth and survival of larval *E. coioides* were associated with the fatty acid composition of the larvae. Larvae with high levels of total fatty acids grew faster than those with lower levels, and larvae with high DHA or EPA content (>2 mg/g DW) exhibited better survival (>10%) than those with lower HUFA content.

Ruangpanit *et al.* (1993) reported improved survival of *E. malabaricus* (*E. coioides*?) larvae when brine shrimp used for larval rearing were enriched with a cod liver oil / egg yolk emulsion ($n=3$ HUFA = 450 mg/g) or were supplemented with copepods or the freshwater cladoceran *Moina*. Dhert *et al.* (1991) found that supplementation of brine shrimp with an emulsion containing high levels of DHA had little effect on growth or survival of *E. coioides* larvae, and concluded that application of DHA could be delayed until D25 before mortality due to nutritional deficiency affected *E. coioides* larvae.

Thyroid hormone treatment

Lam (1994) and Lam *et al.* (1994) reported that the levels of the thyroid hormone are higher in buoyant than in non-buoyant *E. coioides* eggs. The buoyant eggs are more viable than non-buoyant eggs, suggesting a relationship between levels of thyroid hormone and viability. Application of 0.01–1.0 ppm triiodothyronine (T3) or thyroxine (T4) is effective in increasing larval survival and hastening the resorption of the dorsal and anal fins in 2–6 days in contrast to 2–3 weeks in fish not so treated (Tay *et al.*, 1994; de Jesus, 1996; de Jesus *et al.*, 1998). Thyroid hormones can be applied by immersing eggs or larvae in a bath, or by bioencapsulation using rotifers or brine shrimp. De Jesus *et al.* (1998) concluded that a dose rate of 0.01 ppm T4 is appropriate for accelerated metamorphosis and improved survival in 3- to 4-week old grouper larvae.

Commercial grouper larviculture in Taiwan

The only commercial hatchery production of grouper fingerlings that has been identified is from Taiwan. Larviculture of groupers (as well as other marine finfish species) in Taiwan is undertaken

using either the 'indoor method' or 'outdoor method', i.e. in concrete tanks indoors or in outside ponds (Rimmer, 1998).

Indoor method

The indoor method utilises large fibreglass or concrete tanks up to about 100 m³. The rearing tanks are circular or rectangular in shape, flat-bottomed and with a white or light-coloured interior. Larviculture is undertaken using either greenwater or clearwater techniques. The algal density used for greenwater culture ranges from 50,000 to 500,000 cells/ml. Variables such as algal density are only measured in research hatcheries—commercial hatcheries just add algal cells until the desired shade of green is reached (Rimmer, 1998).

Generally, eggs are added directly to the larval rearing tanks. Grouper larvae are fed oyster trochophores from first feed (usually D4) for 2 days. Rotifers are also added to the rearing tanks, generally commencing from first feed. Recent research at Taiwan Fisheries Research Institute's Tungkang Marine Laboratory (TML) indicates that a combination of oyster trochophores and small rotifers (either SS-strain, sieved S-strain, or neonates) is the best initial feed (Su *et al.*, 1997). Rotifer densities are maintained at about 2–3/ml until the grouper larvae are large enough to eat brine shrimp or adult copepods, which is when the dorsal and pectoral spines reach the end of the caudal fin. Generally, grouper larvae are able to feed on adult copepods from D16 (water temperature >26°C) or D22 (<26°C) (Rimmer, 1998).

Outdoor method

The outdoor method of larval rearing is undertaken in concrete or earthen ponds ranging in size from 200 m² to 0.5 ha, and, less commonly, to 1 ha. The ponds are filled only 1–2 days before they are stocked with eggs. The inlet is screened with a fine mesh 'sock' filter to exclude potential predators and nuisance species. Stocking density for grouper ranges from about 1 kg of eggs (i.e. c. 1.5 million eggs) in 0.1 ha to 2 kg (c. 3 million eggs) in 0.2–0.5 ha larval rearing ponds (Rimmer, 1998).

One or two enclosures formed by a tarpaulin set around a floating support structure are set up in the pond, usually with shade cloth overhead to reduce light intensity, and mild aeration to ensure adequate dissolved oxygen and mixing of the water within the enclosure. The enclosures range in size from 5 m³ in small concrete ponds, to 8–10 m³ in earthen ponds 0.2–0.5 ha in area. The enclosures are pumped full of pond water, and fertilised grouper eggs are added to the enclosures. Oyster trochophores are added to the enclosures

from first feed (usually D4) for 2 days, then the larvae are released into the pond. The enclosures allow smaller quantities of oyster trochophores to be fed while retaining relatively high prey densities. They also allow the farmer to visually estimate larval survival after the first few days of culture, which is the period when most mortality occurs. If survival is very low, the farmer may choose to restock the enclosure with another batch of larvae, rather than release the survivors into the pond (Rimmer, 1998).

Rotifers (and, incidentally, other zooplankters) are cultured in small concrete or earthen ponds, usually about 0.05–0.1 ha. The rotifers are cultured using trash fish placed in fertiliser bags and left to decompose in a corner of the pond, or by the addition of organic wastes. A paddle-wheel aerator is placed in the pond to assist with aeration and to generate a current in the pond. Zooplankton is harvested using a fine (c. 85 µm) mesh net that is set downstream from the paddle-wheel aerator for 1–2 hours. The collected zooplankton is added to the larval rearing pond. Farmers attempt to maintain rotifer density at about 3–4/ml, but, as is the case with other aspects of pond management, rotifer density is not measured, but is maintained 'by eye'. Later in the larval rearing cycle, adult copepods are harvested from the zooplankton production ponds using the same technique, although with a larger mesh (c. 210 µm) mesh net. Some farms pump water from the rotifer production ponds into the larval rearing pond, and may also pump zooplankton-rich water from grow-out ponds into the larval rearing ponds (Rimmer, 1998).

The larvae are reared in the ponds until they reach 2.5–3 cm total length (TL), when they are harvested. In the case of grouper, this takes about 4 weeks. Pond temperatures need to be above 20°C to ensure any larval survival for grouper; if pond temperatures drop below 18°C, the grouper larvae will die. For this reason, some farms will not purchase grouper larvae until April, even though eggs are available from early March. In addition, farmers feel that the quality of eggs produced early in the season is inferior to that of eggs produced later in the season. Survival of groupers using both indoor and outdoor larval rearing methods is highly erratic, but generally low: 7 per cent survival is regarded as good. Researchers at TML report that a major constraint to grouper aquaculture is the irregular nature of larval survival. The major problem, according to TML researchers, is high mortality at first feed, although there is often low-level mortality throughout the larval rearing process (Rimmer, 1998). Because of the generally low survival of groupers in larviculture, prices for grouper fingerlings are relatively high: US\$ 2–3 per fingerling (Tamaru *et al.*, 1995).

References

- ASLIANTI, T. (1996). Larval rearing of the grouper *Cromileptes altivelis* at different stocking densities. *Jurnal Penelitian Perikanan Indonesia* 2(2), 6–12. (In Indonesian, English abstract).
- ASLIANTI, T., WARDOYO, J.H. HUTAPEA, S. ISMI & K.M. SETIAWATI. (1998). The polka dots grouper *Cromileptes altivelis* larval rearing under different colours of tank. *Jurnal Penelitian Perikanan Pantai* 4(3), 25–31. (In Indonesian, English abstract).
- CABEROY, N.B. & G.F. QUINTIO. (1998). Sensitivity of grouper *Epinephelus coioides* eggs to handling stress at different stages of embryonic development. *The Israeli Journal of Aquaculture – Bamidgeh* 50, 167–173.
- CHAO, T.M. & L.C. LIM. (1991). Recent developments in the breeding of grouper (*Epinephelus* spp.) in Singapore. *Singapore Journal of Primary Industries* 19, 78–93.
- COWEY, C.B. & J.R. SARGENT. (1972). Fish Nutrition. *Advances in Marine Biology* 10, 383–492.
- DE JESUS, E.G. (1996). Do thyroid hormones play a role in the metamorphosis of the grouper, *Epinephelus coioides*? In: J. Joss (ed.). *The Third Congress of the Asia and Oceania Society for Comparative Endocrinology*, 22–26 January 1991, pp. 259–260. Macquarie University, Sydney, Australia.
- DE JESUS, E.G., J.D. TOLEDO & M.S. SIMPAS. (1998). Thyroid hormones promote early metamorphosis in grouper (*Epinephelus coioides*) larvae. *General and Comparative Endocrinology* 112, 10–16.
- DHERT, P., L.C. LIM, P. LAVEAS, T.M. CHAO, R. CHOU, & P. SORGELOOS. (1991). Effect of dietary essential fatty acids on egg quality and larviculture success of the greasy grouper (*Epinephelus tauvina*, F.): preliminary results. In: P. Lavens, P. Songeloos, E. Jaspers and F. Elsevier (eds). 'Fish and Crustacean Larviculture Symposium'. 27–30 August 1991. Ghent, Belgium. 58–62.
- DOI, M., J. TOLEDO, M.S.N. GOLEZ, M. DE LOS SANTOS & A. OHNO. (1997). Preliminary investigation of feeding performance of larvae of early red-spotted grouper, *Epinephelus coioides*, reared with mixed zooplankton. *Hydrobiologia* 358, 259–263.
- DURAY, M.M. (1994). Daily rates of ingestion of rotifers and artemia nauplii by laboratory-

- reared grouper larvae *Epinephelus suillus*. Philippines Science 31, 32-41.
- DURAY, M.M., C.B. ESTUDILLO & L.G. ALPASAN. (1996). The effect of background colour and rotifer density on rotifer intake, growth and survival of the grouper (*Epinephelus suillus*) larvae. Aquaculture 146, 217-225.
- DURAY, M.M., C.B. ESTUDILLO & L.G. ALPASAN. (1997). Larval rearing of the grouper *Epinephelus suillus* under laboratory conditions. Aquaculture 150, 63-76.
- HUSSAIN, N.A. & M. HIGUCHI. (1980). Larval rearing and development of the brown spotted grouper, *Epinephelus tauvina* (Forskål). Aquaculture 19, 339-350.
- KITAJIMA, C., M. TAKAYA, Y. TSUKASHIMA & T. ARAKAWA. (1991). Development of eggs, larvae and juveniles of the grouper, *Epinephelus septemfasciatus* reared in the laboratory. Japanese Journal of Ichthyology 38, 47-53.
- KOHNO, H., S. DIANI, P. SUNYOTO, B. SLAMET & P.T. IMANTO. (1990). Early developmental events associated with changeover of nutrient sources in the grouper, *Epinephelus fuscoguttatus*, larvae. Bulletin Penelitian Perikanan (Fisheries Research Bulletin) Special Edition No.1, 51-64.
- KOHNO, H., S. DIANI & A. SUPRIATNA. (1993). Morphological development of larval and juvenile grouper, *Epinephelus fuscoguttatus*. Japanese Journal of Ichthyology 40, 307-316.
- KOHNO, H., R.S. ORDONIO-AGUILAR, A. OHNO & Y. TAKI. (1997). Why is grouper larval rearing difficult?: an approach from the development of the feeding apparatus in early stage larvae of the grouper, *Epinephelus coioides*. Ichthyological Research 44, 267-274.
- KUNGVANKIJ, P., L.B. TIRO, B.P. PUDADERA & I.O. POTESTAS. (1986). Induced spawning and larval rearing of grouper (*Epinephelus salmoides*, Maxwell). In: J.L. Maclean, L.B. Dizon & L.V. Hosillos (eds). Proceedings of the First Asian Fisheries Forum, 663-666. Asian Fisheries Society, Manila, Philippines.
- LAM, T.J. (1994). Hormones and egg/larval quality in fish. Journal of the World Aquaculture Society 25, 2-12.
- LAM, T.J., T.M. CHAO, L.C. LIM, D. NUGEGODA & A.M. YONG. (1994). Thyroid hormone levels are higher in buoyant than in non-buoyant eggs in greasy grouper, *Epinephelus tauvina*. Singapore Journal of Primary Industries 22, 29-33.
- LEONG, T.S. (1998). Chapter 13 - Grouper Culture. In: S.S. De Silva (ed.). 'Tropical Mariculture'. Academic Press, London. 423-448.
- LIM, L.C. (1993). Larviculture of the greasy grouper *Epinephelus tauvina* F. and the brown-marbled grouper *E. fuscoguttatus* F. in Singapore. Journal of the World Aquaculture Society 24, 262-274.
- LIM, K.J., C.L. YEN, T.S. HUANG, C.Y. LIM & C.L. CHEN. (1986). Experiment of fry nursing of *E. salmonoides* and its morphological study. Bulletin of the Taiwan Fisheries Research Institute 40, 219-240.
- ORDONIO-AGUILAR, R., H. KOHNO, A. OHNO, M. MOTTEKI & Y. TAKI. (1995). Development of grouper, *Epinephelus coioides*, larvae during changeover of energy sources. Journal of Tokyo University of Fisheries 82, 103-108.
- PECHMANEE, T. & M. ASSAVAAREE. (1993). Nutritional value of rotifer, *Brachionus plicatilis*, fed with emulsified oils rich in n-3 HUFA. In: The Proceedings of Grouper Culture, November 30 - December 1, Songkhla, Thailand. 63-67. National Institute of Coastal Aquaculture, Department of Fisheries, Thailand, and Japan International Cooperation Agency.
- PECHMANEE, T., M. ASSAVAAREE, P. BUNLIPTANON & P. AKKAYANON. (1988). Possibility of using rotifer, *Brachionus plicatilis*, as food for early stage of grouper larvae, *Epinephelus malabaricus*. Technical Paper No. 4. National Institute of Coastal Aquaculture, Department of Fisheries, Thailand. (In Thai.)
- PECHMANEE, T., P. SOMAUEB, M. ASSAVAAREE & S. BOONCHUAY. (1993). The amount of n-3 HUFA in *Chlorella* sp. and *Tetraselmis* sp. In: The Proceedings of Grouper Culture, November 30 - December 1, Songkhla, Thailand. 60-62. National Institute of Coastal Aquaculture and Japan International Cooperation Agency.
- PREDALUMPABURT, Y. & D. TANVILAI. (1988). Morphological development and the early life history of grouper, *Epinephelus malabaricus*, Bloch and Schneider (Pisces: Serranidae). Report of Thailand and Japan Joint Coastal Aquaculture Research Project 3. National Institute of Coastal Aquaculture, Songkhla, Thailand. 49-72.

- QUINITIO, G.F. (1996). The status of grouper and other coral reef fishes seed production in the Philippines. Paper presented at Workshop on Aquaculture of Coral Fishes and Sustainable Reef Fisheries, 4–8 December. Kota Kinabalu, Sabah, Malaysia.
- RIMMER, M.A. (1998). Grouper and snapper aquaculture in Taiwan. *Austasia Aquaculture* 12(1), 3–7.
- RIMMER, M.A., K.C. WILLIAMS & M.J. PHILLIPS. (in press). Proceedings of the Grouper Aquaculture Workshop held in Bangkok, Thailand, 7–8 April 1998.
- RUANGPANIT, N. (1993). Technical manual for seed production of grouper (*Epinephelus malabaricus*). National Institute of Coastal Aquaculture, Department of Fisheries, Ministry of Agriculture and Cooperative, and The Japan International Cooperation Agency. 46 p.
- RUANGPANIT, N., P. BOONLIPTANON & J. KONGKUMNERD. (1993). Progress in the proportion and larval rearing of the grouper, *Epinephelus malabaricus*. In: The Proceedings of Grouper Culture, 30 November – December 1, 1993. National Institute of Coastal Aquaculture, Songkhla, Thailand and Japan International Cooperation Agency.
- SU, H.M., M.S. SU & I.C. LIAO. (1997). Preliminary results of providing various combinations of live foods to grouper (*Epinephelus coioides*) larvae. *Hydrobiologia* 358, 301–304.
- SU, H.M., M.S. SU & I.C. LIAO. (1998). The culture and use of microalgae for larvae rearing in Taiwan. In: Abstracts of the Joint Taiwan–Australia Aquaculture and Fisheries Resources and Management Forum, held at Keelung, Taiwan, 2–8 November 1998. p. 28.
- TAMARU, C.S., F. CHOLIK, J.C-M. KUO & J.JR. FITZGERALD. (1995). Status of the culture of milkfish (*Chanos chanos*), striped mullet (*Mugil cephalus*) and grouper (*Epinephelus* sp.). *Reviews in Fisheries Science* 3(3), 249–273.
- TAY, H.C., J. GOH, A.N. YONG, H.S. LIM, T.M. CHAO, R. CHOU *et al.* (1994). Effect of thyroid hormone on metamorphosis in greasy grouper, *Epinephelus tauvina*. *Singapore Journal of Primary Industries* 22, 35–38.
- TOLEDO, J.D., S.N. GOLEZ, M. DOI & A. OHNO. (1997). Food selection of early grouper, *Epinephelus coioides*, larvae reared by the semi-intensive method. *Suisanzoshoku* 45, 327–337.
- TOLEDO, J.D., S.N. GOLEZ, M. DOI & A. OHNO. (1999). Use of copepod nauplii during early feeding stage of grouper *Epinephelus coioides*. *Fisheries Science* 65, 390–397.
- WARDOYO, K.M. SETIAWATI, S. ISMI, J. HUTAPEA & T. ASLIANTI. (1997). Effect of rotifer density on growth and survival rate of polka dots grouper larvae (*Cromileptes altivelis*). In: *Prosiding Simposium Perikanan Indonesia II, Ujung Pandang, 2–3 December 1997*. 234–237. (In Indonesian, English abstract).
- WASPADA, MAYUNAR & T. FATONI. (1991a). Nutrient enrichment of rotifers, *Brachionus plicatilis*, to support fry production of grouper, *Epinephelus fuscoguttatus*. *Jurnal Penelitian Budidaya Pantai* 7, 73–79. (In Indonesian, English abstract).
- WASPADA, S. MURTININGSIH, T. AHMAD & M. SUHARJO. (1991b). Effect of different tank shapes on the growth and survival rate of grouper, *Epinephelus fuscoguttatus* larvae. *Jurnal Penelitian Budidaya Pantai* 7, 42–50. (In Indonesian, English abstract).
- WASPADA, Y. SETIAWAN & M. RODIF. (1991c). Effect of enriched rotifers, *Brachionus plicatilis* on the growth and survival rate of grouper, *Epinephelus fuscoguttatus*, larvae. *Jurnal Penelitian Budidaya Pantai* 7, 57–66. (In Indonesian, English abstract).
- WATANABE, W.O., S.C. ELLIS, E.P. ELLIS, V.G. LOPEZ, P. BASS, J. GINOZA & A. MORIWAKE. (1996). Evaluation of first-feeding regimens for larval Nassau grouper *Epinephelus striatus* and preliminary pilot-scale culture through metamorphosis. *Journal of the World Aquaculture Society* 27, 323–331.

